



ABOUT METASTATIC HER2+ BREAST CANCER

What is HER2+ breast cancer?

HER2+ (human epidermal growth factor receptor 2-positive) breast cancer is different from other types of breast cancer. It tends to be more aggressive, and its chance of returning or progressing is a reality—even after treatment with other regimens. In some breast cancers, cancer cells have a gene mutation that makes too much HER2 protein. This protein promotes the growth of cancer cells. Among women with breast cancer, approximately 20% have HER2+ breast cancer. HER2+ breast cancer can grow fast and has a high chance of coming back and spreading in the body.

What is metastatic HER2+ breast cancer?

Metastatic breast cancer (mBC)—also called Stage 4 or advanced breast cancer—is cancer that has spread beyond the breast and nearby lymph nodes to other parts of the body. Treatment can control metastatic breast cancer for some time, but often the cancer continues to grow or spread. A person's experience with metastatic breast cancer will depend on how many metastases develop and where they are in the body.



ABOUT NERLYNX

What is NERLYNX (ner links)?

NERLYNX is a HER2-targeted small molecule medication approved by the U.S Food and Drug Administration (FDA) for metastatic HER2+ breast cancer.

Who is eligible to take NERLYNX?

NERLYNX is used with a medicine called capecitabine to treat adults with HER2+ breast cancer that has spread to other parts of the body (metastatic) **and** who have received 2 or more anti-HER2 therapy medicines for metastatic breast cancer.

It is not known if NERLYNX is safe and effective in children.

How does NERLYNX work?

In people with breast cancer, cells receive signals that cause the cells to grow and divide. NERLYNX is a type of treatment called a targeted therapy that works inside the cell to block these signals that tell the cells to grow and divide. By blocking HER2, breast cancer cells do not grow and divide as rapidly.

How effective is NERLYNX?

NERLYNX was studied in a clinical trial of 621 people who had HER2+ mBC and had undergone treatment with 2 or more HER2-directed treatments for mBC. The phase 3 study examined the effectiveness of NERLYNX plus capecitabine. The study compared people who took NERLYNX plus capecitabine (a chemotherapy) with people who took lapatinib (Tykerb[®]) plus capecitabine. In the clinical trial, NERLYNX reduced the risk of cancer progressing by 24% for all people studied who had undergone treatment with 2 or more HER2-directed treatments for mBC. People taking NERLYNX responded to treatment longer than those taking lapatinib (median of 8.5 months compared to 5.6 months for those taking lapatinib [Tykerb[®]]).*

What are the most common side effects of NERLYNX?

The most common side effects of NERLYNX in combination with capecitabine include: diarrhea, nausea, vomiting, decreased appetite, constipation, tiredness/weakness, weight loss, dizziness, back pain, joint pain, urinary tract infection, upper respiratory tract infection, swelling of your stomach-area, kidney problems, and muscle spasms.

These are not all of the possible side effects of NERLYNX. For more information, ask your Healthcare Provider.

Tell your doctor if you have any side effects that bother you or that does not go away. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

*The 8.5 and 5.6 months durations were calculated only from the percentage of people who responded to treatment (32.8% in the NERLYNX group vs 26.7% in the lapatinib group).

WHAT IS NERLYNX?

NERLYNX is also used with a medicine called capecitabine to treat adults with HER2-positive breast cancer that has spread to other parts of the body (metastatic) and who have received 2 or more anti-HER2 therapy medicines for metastatic breast cancer.

It is not known if NERLYNX is safe and effective in children.

For additional IMPORTANT SAFETY INFORMATION, including [Patient Information](#), please see pages 4 through 6 of this PDF.



BEFORE TAKING NERLYNX

Do I need to be re-tested for HER2 before taking NERLYNX?

If you have already tested positive for HER2+ breast cancer, you should ask your healthcare team whether you need to be retested before taking NERLYNX. However, in order to receive NERLYNX for mBC, you must have received 2 or more anti-HER2 therapy medicines for metastatic breast cancer.

What if I'm not HER2+? Can I still take NERLYNX?

NERLYNX is a treatment for HER2+ breast cancer patients only. Talk to your healthcare team about the treatment options that may be right for you.

Is NERLYNX covered by insurance?

NERLYNX is covered by many insurance providers. For questions about insurance, the Puma Patient Lynx™ support program has resources for you to explore. Please see the next page for additional program details.

Where can I get NERLYNX?

Your doctor can provide you with a prescription that is filled through a network of specialty pharmacies. If you have questions related to your specific situation, please speak with your healthcare team.



WHILE TAKING NERLYNX

How can I best manage side effects I may experience?

As with most medicines, side effects can occur with NERLYNX. Working with your healthcare team from the beginning is the key to success. Be sure to let your doctor know about any side effects that you experience and ask for help to manage them.

Your doctor can prescribe medication, recommend over-the-counter medicines, or reduce your dose of NERLYNX to help. Diarrhea is the most common side effect. You will take NERLYNX with an antidiarrheal, as needed. Your doctor may start you on the full dose of 6 tablets of NERLYNX and then reduce the dose as needed. These reductions can help manage side effects while still allowing NERLYNX to work for you.

How do I take NERLYNX and for how long?

NERLYNX is an oral medicine that you take once a day. You will take NERLYNX in combination with capecitabine (a chemotherapy). **For 14 days**, you will take 6 NERLYNX tablets once a day plus capecitabine twice a day. After 14 days, you will stop taking capecitabine for 7 days. **You will continue taking NERLYNX during this time.** After 7 days, you will add capecitabine back again for 14 days. The cycle keeps repeating.

You will also need to take loperamide, an antidiarrheal, during the first 2 months of NERLYNX treatment and then as needed. Talk to your doctor about how much and how often to take this medicine.

What if I miss a dose of NERLYNX?

If you miss a dose of NERLYNX, skip that dose and take your next dose at your regular scheduled time. If you have questions related to your specific situation, please speak with your healthcare team.

SELECT IMPORTANT SAFETY INFORMATION

What is the most important information I should know about NERLYNX (ner links)?

NERLYNX® (neratinib) may cause serious side effects, including:

- **Diarrhea.** Diarrhea is a common side effect of NERLYNX, but it can also be severe. You may lose too much body salts and fluid, and get dehydrated. When you start NERLYNX, your healthcare provider should prescribe the medicine loperamide for you during your first 2 months (56 days) of NERLYNX and then as needed. Be sure that your healthcare provider prescribes anti-diarrheals with NERLYNX. Anti-diarrheals must be started with the first dose of NERLYNX.

For additional IMPORTANT SAFETY INFORMATION, including [Patient Information](#), please see pages 4 through 6 of this PDF.



PUMA PATIENT LYNX™ SUPPORT PROGRAM

What services are available to support patients interested in or currently taking NERLYNX?

Puma Patient Lynx programs and services provide ongoing education, encouragement, and financial assistance:

Ongoing Support

NERLYNX Mentor Program: When you start taking NERLYNX, the mentor program connects you with someone who currently takes or has taken NERLYNX.

Nurse Support: Nurses are available, Monday through Friday 9 AM - 8 PM EST, to answer your NERLYNX questions through the call center.

Text Support Program: Receive guidance and reassurance through emails and texts that will help you stay committed to taking NERLYNX, remember to take your daily medicine, manage side effects, reduce stress and anxiety, and get the most out of your healthcare appointments.

Access Programs

NERLYNX Quick Start: Provides a 3-week supply of NERLYNX to eligible patients whose access to medicine is delayed.

Patient Assistance Programs: For information or assistance with pharmacy insurance benefits, call 1-855-816-5421.

Co-pay Support: Commercially insured, eligible patients treated with NERLYNX may pay as little as \$10 per prescription.[†]

Supportive Care Voucher: Patients can receive up to a 3-month supply of products used for antidiarrheal treatment for free.[‡]

Specialty Pharmacy Services: NERLYNX is available through specialty pharmacies that help with much more than filling your prescription. They provide insurance verification, help with prior authorization requirements for insurance, on-time delivery of NERLYNX, nurses, and pharmacist counselors to help with questions about NERLYNX and side effects.

To learn more about the Puma Patient Lynx support program, visit [NERLYNX.com](https://www.nerlynx.com) or call 1-855-816-5421, Monday through Friday, 9 AM - 8 PM EST, to connect with someone for more information.

YOU CAN ALSO LEARN MORE ABOUT NERLYNX
BY CONNECTING WITH US



[†]Limitations apply. Patient must have commercial insurance. Offer is not valid under Medicare, Medicaid, or any other federal or state program. Puma Biotechnology reserves the right to rescind, revoke, or amend this program without notice. For full terms and conditions, call 1-855-816-5421.

[‡]This program benefit is offered based on the findings from the CONTROL trial, which show that using antidiarrheals can help to manage diarrhea that may be caused by NERLYNX. Limitations apply. Puma Biotechnology reserves the right to rescind, revoke, or amend this program without notice. For full terms and conditions, visit www.nerlynx.com or call 1-855-816-5421.

IMPORTANT SAFETY INFORMATION

What is the most important information I should know about NERLYNX (ner links)?

NERLYNX® (neratinib) may cause serious side effects, including:

- **Diarrhea.** Diarrhea is a common side effect of NERLYNX, but it can also be severe. You may lose too much body salts and fluid, and get dehydrated. When you start NERLYNX, your healthcare provider should prescribe the medicine loperamide for you during your first 2 months (56 days) of NERLYNX and then as needed. Be sure that your healthcare provider prescribes anti-diarrheals with NERLYNX. Anti-diarrheals must be started with the first dose of NERLYNX. To help prevent or reduce diarrhea:
 - You should start taking loperamide with your first dose of NERLYNX.
 - Keep taking loperamide during the first 2 months (56 days) of NERLYNX treatment and then as needed. Your healthcare provider will tell you exactly how much and how often to take this medicine.
 - Your healthcare provider may also need to give you other medicines to manage diarrhea when you start treatment with NERLYNX. Follow your healthcare provider's instructions on how to use these anti-diarrheal medicines.
 - Always take anti-diarrheals exactly as your healthcare provider tells you.
 - While taking anti-diarrheals, you and your healthcare provider should try to keep the number of bowel movements that you have at 1 or 2 bowel movements each day.
 - Tell your healthcare provider if you have more than 2 bowel movements in 1 day, or you have diarrhea that does not go away.
 - **Call your healthcare provider right away, as instructed, if you have severe diarrhea or if you have diarrhea along with weakness, dizziness, or fever.**
 - After 2 months (56 days) of treatment with NERLYNX, follow your healthcare provider's instructions about taking loperamide as needed to control diarrhea.

Your healthcare provider may change your dose of NERLYNX, temporarily stop or completely stop NERLYNX if needed to manage your diarrhea.

See "What are the possible side effects of NERLYNX?" for more information about side effects.

What is NERLYNX?

- NERLYNX is also used with a medicine called capecitabine to treat adults with HER2-positive breast cancer that has spread to other parts of the body (metastatic) **and** who have received 2 or more anti-HER2 therapy medicines for metastatic breast cancer.

It is not known if NERLYNX is safe and effective in children.

Before taking NERLYNX, tell your healthcare provider about all of your medical conditions, including if you:

- have liver problems. You may need a lower dose of NERLYNX.
- are pregnant or plan to become pregnant. NERLYNX can harm your unborn baby. If you are a female who can become pregnant:
 - Your healthcare provider should do a pregnancy test before you start taking NERLYNX.
 - You should use effective birth control (contraception) during treatment and for at least 1 month after your last dose of NERLYNX.
 - Talk with your healthcare provider about forms of birth control that you can use during this time.
 - Tell your healthcare provider right away if you become pregnant during treatment with NERLYNX.
 - Males with female partners who can become pregnant should use effective birth control during treatment and for 3 months after your last dose of NERLYNX.
- are breastfeeding or plan to breastfeed. It is not known if NERLYNX passes into your breast milk. Do not breastfeed during treatment and for at least 1 month after your last dose of NERLYNX.

Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements. Especially tell your healthcare provider if you take medicines used to decrease stomach acid, called proton pump inhibitors or PPIs. You should avoid taking these medicines during treatment with NERLYNX.

What are the possible side effects of NERLYNX?

NERLYNX may cause serious side effects, including:

See "What is the most important information I should know about NERLYNX?"

- **Liver problems.** Changes in liver function tests are common with NERLYNX. Your healthcare provider should do blood tests before you begin treatment, monthly during the first 3 months, and then every 3 months as needed during treatment with NERLYNX. Your healthcare provider will stop your treatment with NERLYNX if your liver tests show severe problems. Call your healthcare provider right away if you get any of the following signs or symptoms of liver problems:

- | | |
|--|---|
| - tiredness | - fever |
| - nausea | - rash |
| - vomiting | - itching |
| - pain in the right upper stomach-area (abdomen) | - yellowing of your skin or whites of your eyes |

The most common side effects of NERLYNX when used with capecitabine include:

- | | |
|----------------------|-------------------------------------|
| · diarrhea | · back pain |
| · nausea | · joint pain |
| · vomiting | · urinary tract infection |
| · decreased appetite | · upper respiratory tract infection |
| · constipation | · swelling of your stomach-area |
| · tiredness/weakness | · kidney problems |
| · weight loss | · muscle spasms |
| · dizziness | |

These are not all of the possible side effects of NERLYNX. For more information, ask your Healthcare Provider.

Tell your doctor if you have any side effects that bother you or that does not go away. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

For [Patient Information](#), please see pages 5 and 6.



NERLYNX is a registered trademark and Puma Patient Lynx is a trademark of Puma Biotechnology, Inc.

Tykerb is a registered trademark of Novartis, Inc.

© 2020 Puma Biotechnology, Inc. All Rights Reserved. PRC-US-NER-1725 08/20

nerlynx[®]
(neratinib) tablets

PATIENT INFORMATION

NERLYNX® (ner links)

(neratinib)

tablets

What is the most important information I should know about NERLYNX?

NERLYNX may cause serious side effects, including:

- **Diarrhea.** Diarrhea is a common side effect of NERLYNX, but it can also be severe. You may lose too much body salts and fluid, and get dehydrated. When you start NERLYNX, your healthcare provider should prescribe the medicine loperamide for you during your first 2 months (56 days) of NERLYNX and then as needed. Be sure that your healthcare provider prescribes anti-diarrheals with NERLYNX. Anti-diarrheals must be started with the first dose of NERLYNX. To help prevent or reduce diarrhea:
 - You should start taking loperamide with your first dose of NERLYNX.
 - Keep taking loperamide during the first 2 months (56 days) of NERLYNX treatment and then as needed. Your healthcare provider will tell you exactly how much and how often to take this medicine.
 - Your healthcare provider may also need to give you other medicines to manage diarrhea when you start treatment with NERLYNX. Follow your healthcare provider's instructions on how to use these anti-diarrheal medicines.
 - Always take anti-diarrheals exactly as your healthcare provider tells you.
 - While taking anti-diarrheals, you and your healthcare provider should try to keep the number of bowel movements that you have at 1 or 2 bowel movements each day.
 - Tell your healthcare provider if you have more than 2 bowel movements in 1 day, or you have diarrhea that does not go away.
 - **Call your healthcare provider right away, as instructed, if you have severe diarrhea or if you have diarrhea along with weakness, dizziness, or fever.**
 - After 2 months (56 days) of treatment with NERLYNX, follow your healthcare provider's instructions about taking loperamide as needed to control diarrhea.

Your healthcare provider may change your dose of NERLYNX, temporarily stop or completely stop NERLYNX if needed to manage your diarrhea.

See "**What are the possible side effects of NERLYNX?**" for more information about side effects.

What is NERLYNX?

- NERLYNX is a prescription medicine used alone to treat adults with early-stage human epidermal growth factor receptor 2 (HER2)-positive breast cancer **and** who have previously been treated with trastuzumab-based therapy.
- NERLYNX is also used with a medicine called capecitabine to treat adults with HER2-positive breast cancer that has spread to other parts of the body (metastatic) **and** who have received 2 or more anti-HER2 therapy medicines for metastatic breast cancer.

It is not known if NERLYNX is safe and effective in children.

Before taking NERLYNX, tell your healthcare provider about all of your medical conditions, including if you:

- have liver problems. You may need a lower dose of NERLYNX.
- are pregnant or plan to become pregnant. NERLYNX can harm your unborn baby. If you are a female who can become pregnant:
 - Your healthcare provider should do a pregnancy test before you start taking NERLYNX.
 - You should use effective birth control (contraception) during treatment and for at least 1 month after your last dose of NERLYNX.
 - Talk with your healthcare provider about forms of birth control that you can use during this time.
 - Tell your healthcare provider right away if you become pregnant during treatment with NERLYNX.
 - Males with female partners who can become pregnant should use effective birth control during treatment and for 3 months after the last dose of NERLYNX.
- are breastfeeding or plan to breastfeed. It is not known if NERLYNX passes into your breast milk. Do not breastfeed during treatment and for at least 1 month after your last dose of NERLYNX.

Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements. Especially tell your healthcare provider if you take medicines used to decrease stomach acid, called proton pump inhibitors or PPIs. You should avoid taking these medicines during treatment with NERLYNX.

How should I take NERLYNX?

- Take NERLYNX exactly as your healthcare provider tells you to take it.
- Your healthcare provider may change your dose of NERLYNX if needed.
- Take NERLYNX with food.

- Take NERLYNX at about the same time each day.
- Swallow NERLYNX tablets whole. Do not chew, crush, or split NERLYNX tablets.
- If you take an antacid medicine, take NERLYNX 3 hours after the antacid medicine.
- If you take an acid reducer (H₂ receptor blocker), NERLYNX should be taken at least 2 hours before or 10 hours after you take these medicines.
- NERLYNX is usually taken for 1 year.
- If you miss a dose of NERLYNX, skip that dose and take your next dose at your regular scheduled time.
- If you take too much NERLYNX, call your healthcare provider right away or go to the nearest hospital emergency room.

What should I avoid while taking NERLYNX?

You should avoid eating products that contain grapefruit during treatment with NERLYNX.

What are the possible side effects of NERLYNX?

NERLYNX may cause serious side effects, including:

See “What is the most important information I should know about NERLYNX?”

- **Liver problems.** Changes in liver function tests are common with NERLYNX. Your healthcare provider should do blood tests before you begin treatment, monthly during the first 3 months, and then every 3 months as needed during treatment with NERLYNX. Your healthcare provider will stop your treatment with NERLYNX if your liver tests show severe problems. Call your healthcare provider right away if you get any of the following signs or symptoms of liver problems:
 - tiredness
 - nausea
 - vomiting
 - pain in the right upper stomach-area (abdomen)
 - fever
 - rash
 - itching
 - yellowing of your skin or whites of your eyes

The most common side effects of NERLYNX when used alone include:

- diarrhea
- nausea
- stomach-area (abdomen) pain
- tiredness
- vomiting
- rash
- dry or inflamed mouth, or mouth sores
- decreased appetite
- muscle spasms
- upset stomach
- nail problems including color change
- dry skin
- swelling of your stomach-area
- nosebleed
- weight loss
- urinary tract infection

The most common side effects of NERLYNX when used with capecitabine include:

- diarrhea
- nausea
- vomiting
- decreased appetite
- constipation
- tiredness/weakness
- weight loss
- dizziness
- back pain
- joint pain
- urinary tract infection
- upper respiratory tract infection
- swelling of your stomach-area
- kidney problems
- muscle spasms

These are not all of the possible side effects of NERLYNX. For more information, ask your healthcare provider. Tell your doctor if you have any side effects that bother you or that does not go away. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

How should I store NERLYNX?

- Store NERLYNX at room temperature between 68°F to 77°F (20°C to 25°C).

Keep NERLYNX and all medicines out of the reach of children.

General information about the safe and effective use of NERLYNX.

Medicines are sometimes prescribed for purposes other than those listed in a Patient Information leaflet. Do not use NERLYNX for a condition for which it was not prescribed. Do not give NERLYNX to other people, even if they have the same symptoms that you have. It may harm them. You can ask your pharmacist or healthcare provider for information about NERLYNX that is written for health professionals.

What are the ingredients in NERLYNX?

Active ingredient: neratinib

Inactive ingredients: Tablet Core: colloidal silicon dioxide, mannitol, microcrystalline cellulose, crospovidone, povidone, magnesium stearate, and purified water. Coating: red film coat: polyvinyl alcohol, titanium dioxide, polyethylene glycol, talc, and iron oxide red.

Manufactured for: Puma Biotechnology, Inc. 10880 Wilshire Blvd., Suite 2150 Los Angeles, CA 90024-4106
©2020, Puma Biotechnology, Inc. All rights reserved.